Advanced Study_Occup Safety

by kresna febriyanto

Submission date: 19-Jun-2023 06:33AM (UTC-0400)

Submission ID: 2118971521

File name: 2nd_Paper.pdf (239.25K)

Word count: 5264 Character count: 26807

Assessment of Occupational Noise Exposure and Hearing Disturbances: A Study among Sp. Speedboat Workers

Kresna Febriyanto^{12*}, Joana Cristina Cardoso Guedes¹

¹Faculty of Engineering, University of Porto, Portugal

²Faculty of Public Health, Universitas Muhammadiyah Kalimantan Timur, Indonesia

*Correspondence author: up202202416@edu.fe.up.pt

ABSTRACT

Introduction: The aims of present study were classify levels of occupational noise, hearing disturbances, and identify other factors that impact hearing loss. Method: From July to August 2022, this cross-sectional study enrolled speedboat members. We used sound level meter as a noise measurement, and audiometer for assessing hearing problems. The questionnaires were used to support the data with the questions according to age, job position, Body Mass Index (BMI), years of working, work duration in a day, smoking and alcohol consumption, physical activities, and hobbies during leisure time. Chi-Square test was used to find out the relationship between the variables. Result: The captain's position does not exceed the threshold limit value (TLV) (average noise level was 85.45 dBA). In contrast, six speedboats in the crew's location exceeded TLV based on the length of exposure duration. A total of 109 participants (83.8%) had hearing loss, with the majority suffering from mild (40.8%), moderate (39.2%), and severe (3.8%) levels. **Conclusion:** Except for job position, all independent variables were related to hearing impairment. Due to prolonged exposure to noise, high decibel levels, and other factors, members of speedboats are susceptible to hearing loss. Age, BMI, and other lifestyle factors such as smoking, alcohol consumption, recreational activities, and daily exercise will also contribute to their hearing loss.

Keywords: Occupational Noise, Hearing Impairment, Vessel, Ship, Seafarers, Sailors

Introduction

Noise-induced hearing loss (NIHL), one of the most prevalent forms of sensorineural hearing loss among individuals working in noisy world's environments, is the most pervasive industrial disease in the world (1). There are three fundamental forms of hearing loss: conductive loss, sensorineural loss, and mixed loss. Once the hair cells in the cochlea are damaged, sensorineural hearing loss is usually permanent (2).

The NIHL has been studied in a variety of industrial and nonindustrial activities, including the transportation industry. In the era of globalization, transport is a crucial need for the community to support all activities and daily rituals. Each mode of transportation has its function and capacity for passenger service. Maritime transport is

the mode of public transport that assists the government in its genuine development efforts. Sea transport is essential for connecting islands so that the distribution of products and passengers from one island to another can run smoothly and development can be distributed equitably rather than being concentrated in a single region or island (3).

Noise exposure in the workplace is one of the primary causes of NIHL. Because they are exposed to noise daily, the sailors and crew members are susceptible to NIHL. There are several sources of noise on a boat, including engines, wind, waves, and entertainment on board. Oldenburg et al. (2020) mention that high levels of noise on board are caused by ship operation, loading and offloading (especially from container ships), port stay, port facilities

(such as cranes), and environmental noise (wind and sea) (4).

Noise pollution is a new emerging environmental issue caused by ships (5). Noise exposure is one of the most prevalent occupational risk factors associated with hearing loss. Hearing impairment can be detected early before permanent hearing loss results from continued exposure (6).

According to the most recent estimates, 1.59 billion persons worldwide, or 20.3% of the global population, are affected by hearing loss, with 430 million (5.5%) suffering from moderate or severe hearing loss (7). By 2050, it is anticipated that the number of persons with hearing loss will approach 2.5 billion, with 700 million requiring intervention. Hearing loss was classified as the third leading cause of years lived with a disability globally in 2019, according to statistics (8). Globally, hearing loss is estimated to cause an annual economic burden of over \$750 billion (9). In a study conducted on five distinct Brazilian vessels, approximately 56.5% of the crew suffered from NIHL. At 78.8%, engine room personnel had the highest incidence of NIHL (10). Such damage to the hearing system makes it challenging to communicate with the encompassing environment comprehend the verbal communication of others, especially in noisy environments. As a result, it can also increase the likelihood of occupational accidents (6,11-13). 22

The International Maritime Organization's (IMO) Maritime Safety Committee, the primary international body regulating safety at sea from hazardous noise pressure levels in vessels, has issued a warning to ship operators: acceptable noise levels are 85 dBA in work environments, 75 dBA in motor control rooms, 65 dBA in command and navigation spaces, and 60 dBA in common areas (14). As for the ship type, the noise threshold value is not expressly stated but Errowas modified according to exposure ing

duration. According to the regulations, employees may be exposed to 85 dBA for no more than eight hours daily reeven though the threshold value for noise on ships has been defined, many research investigations on noise intensity were above the limit.

Previous studies conducted on seafarers only examined hearing loss resulting from shipboard noise. They Error measured the noise level on the vessels and the workers' hearing thresholds. Hearing loss is not caused exclusively by noise but also by non-occupational factors such as age, work experience, body mass index (BMI), and non-infectious chronic diseases

Therefore, this study was conducted to classify levels of occupational noise, hearing disturbances, and identify other factors that impact hearing loss.

Materials and Methods

Study design

The present research was designed as a cross-sectional study to investigate hearing disturbances experienced to noise in operating the determine other factors that correlate to their hearing problems.

Place and time of study

The study was conducted during July to August 2022. Study data were collected within working hours of the day at Tengkayu II, port of Tarakan City. Tarakan is an island in North Borneo province, Indonesia that has a vital role in advancing equitable development in this province. Through Tarakan, it becomes the entrance gate and a transit city connecting North Kalimantan areas.

Study population and sample

There were 168 speedboat workers included in this survey, with 56 speedboats operating at the port and each speedboat containing three to four workers. The researcher used the Slovin formula to

calculate the sample size from N = 168. The calculation formula is as follows:

$$n = \frac{N}{1 + N(d)^2}$$

Where N is the size of the population, and d is the desired level of accuracy (0.05).

The calculation results indicate 118 respondents, but to avoid errors in data acquisition or loss, the researchers added 10%, bringing the total number of samples in this study (n) to 130 respondents. The sampling technique utilized in this study was simple random sampling, specifically sampling of respondents who could be in a location that corresponds with the research context (15).

Data collection tools Noise measurements

Noise measurements in speedboats have been carried out on 12 speedboats. Using the Sound Level Measurement (SLM) type Lutron SL 4001, each speedboat was measured at two locations (the skipper's and the crew's positions) while in operation. The SLM was placed parallel to the worker's ear for two to three minutes to measure the continuous noise generated by the ship's engine that entered the worker's ears. As shown in Figure 1, the type of vessel operating in the Tarakan region is a speedboat with a capacity of 15 to 60 passengers.

Hearing assessment

To determine the air conduction threshold, pure tone audiometry was performed. The subjects were not exposed to occupational noise for at least 14 hours prior to the audiometric threshold testing to recovery from temporary threshold shift. The result of hearing impairment was classified into: normal (< 25 dB), mild (26-40 dB), moderate (41-55 dB), and severe (> 55 dB) (16). They were evaluated in a soundproof chamber using a pure tone

audiometer. Hearing levels at 500, 1,000, 2,000, 4,000, and 8,000 kHz were determined using the ascending threshold technique.

Questionnaire

The questionnaire included queries regarding occupational and non-occupational risk factors for hearing impairment (Table 1). This included questions about personal identification (such as age and BMI), the current job (job position, years of working, work duration in a day), and health habits (smoking and alcohol consumption, exercise activities, hobbies).

Statistical analysis

SPSS (version 22) was used to calculate descriptive statistic (mean, min, max, SD, frequency, and percentages) and bivariate analyzed among the variables. Chi-Square test was used to measure association a between each independent variables (noise level, age, job description, body mass index, smoking consumption, years of working, work duration in a day, drinking alcohol, exercise activities, and hobbies related to loud sound) against dependent variable (hearing disturbance). The significant association was define as p value < 0.05.

Ethical approval

Since the use of humans in research necessitates the protection of individual rights, the "informed consent" condition was implemented as a code of ethics, and each employee's verbal permission was obtained. This research has also passed the health research ethics test. As stated in the Letter of Agreement for Consideration No. 91/KEPK-FK/VII/2022 issued by the Health Research Ethics Commission, Faculty of Medicine. Mulawarman University on July 5, 2022.





Figure 1. The Speedboat

Results

The majority of speedboat employees are between the ages of 36 and 45, with an average age of 37 years old, as indicated by Table 1 as primary characteristics. Despite being classified as the old-aged adult age group, the majority workers have worked less than seven years (51.5%). Most employees have a normal body mass index, with an average of 15 cigarettes smoked daily. However, more than 80% of workers were cigarette smokers. Only 43 respondents did not drink alcohol, and only 21 participated in physical activities outside their profession. Over fifty percent of employees have noise-related hobbies.

According to the findings of interviews, employees engage in noise-related activities such as: went to nightclubs, listening to music through loudspeakers or headphones, concert attendance, and hunting/shooting.

Therefore, each speedboat has between three and four crew members, and the majority of respondents in this study (84.6%) are also speedboat personnel A total of 109 respondents out of 130 fused research samples had hearing loss, the majority of whom had mild (53 respondents) and moderate (51 respondents) levels.

Table 1. Demographic and general information of speedboat workers

Variable	and general in community of Species	Frequency (n=130)	Lercentage
v ariabie			Frag. ((%)
Age	Late tens (17-25)	10	7.7
	Young adults (26-35)	40	30.8
	Old-aged adults (36-45)	58	44.6
	Young elderly (46-55)	21	16.1
	Late elderly (56-65)	1	0.8
	Min: 19; Max: 59; Mean: 30	6.99	
Job position	Skipper	20	15.4
-	Speed boat Crews	110	84.6
Body mass index	Normal weight	74	56.9
	Pre-obesity	3	2.3
	Obesity class I	24	18.5
	Obesity Class II	29	22.3
Smoking consumption	No	19	14.6
	Yes	111	85.4
	Min: 0; Max: 32; Mean: 14.53 (bat	rs in a day)	
Years of Working	≤7 years	67	51.5
	>7 years	63	48.5
	Min: 1; Max: 20; Mean: 7.	42	
Work duration (in a day)	1 – 5 hr	82	63.1
	>5 hr	48	36.9
	Min: 2; Max: 6; Mean: 3.8	86	
Drinking alcohol	No	43	33.1
Missing "	" (ETS) Yes	87	66.9

Variable		Frequency (n=130)	Percentage			
variabie			(%)			
Min: 0; Max: 4; Mean: 1 26						
Physical activities	Yes	21	16.2			
	No	109	83.8			
Time to do exercise	Never do exercise	109	83.8			
Proofread (FIS)	Daily	4	3.1			
110011 cad	Weekly	13	10			
	Monthly	4	3.1			
Hobbies	Not related with high sound	54	41.5			
	Related with high sound	76	58.5			
Hearing status	Normal Prep. (ES)	21	16.2			
Z .	Disturbances	109	83.8			
Hearing Disturbances	Normal	21	16.2			
	Mild	53	40.8			
	Moderate	51	39.2			
	Severe	5	3.8			

Table 2. Results of noise assessment

No	Length of journey	ror Kind of boat	SNumber of Engines	Route	Pos.1 (skipper)	Pos.2 ne (crew) Tone (ETS
1	1.5-2 hr	closed	5 (@200pk)	A - B	85.4	91.6
2	1.5-2 hr	open	5 (@200pk)	B-A	86.9	92.1
2 3 4 5	1.5-2 hr	open	4 (@200pk)	A - B	85.7	87.4
4	1.5-2 hr	closed	4 (@200pk)	B-A	83.9	87.6
5	3.5-4 hr	closed	5 (@200pk)	A - C	87	92
6 7	3.5-4 hr	open	5 (@200pk)	C-A	88.4	90.2
7	3.5-4 hr	closed	4 (@200pk)	A-D	85.6	88.3
8	3.5-4 hr	open	4 (@200pk)	D-A	87.1	88.5
9	1 hr	open Miss	3 (@200pk)	A - E	83.2	86.7
10	1 hr	closed	4 (@200pk)	E - A	81.3	88.1
11	1 hr	open	1 (@200pk)	$\mathbf{A} - \mathbf{E}$		84.2
12	1.5 hr	open	1 (@200pk)	A - B		84.9
		Min			81.3	84.2
		Max			88.4	92.1
		Average			85.45	88.47

issing "," 🙉

The measurements of noise using the SLM were showed in Table 2. The number of engines per speedboat ranges from one to five, with a power of 200pk. Each speedboat typically makes two round journeys per day with travel time one to four hours per way. There were two types of speedboats: closed (speedboat walls cover the entire speedboat body, as shown in Figure 1) and open (speedboat walls cover only half or three-quarters of the ship).

In this study, researchers measured the noise of four speedboat routes (round trip

each routes), centered on the port of Tengkayus H, tarakan City (where A is Tarakan). In accordance with international fused regulations and the decision of the Indonesian Minister of Manpower, the noise threshold limit value (TLV) for four hours of work is 88 dBA; for two hours, it is 91 dBA; and for one hour, it is 94 dBA (17). Noise measurements results showed that overall the skipper position did not exceed the TLV. In contrast, in the crew's place, six speedboats exceeded the threshold value according to the length of exposure (speedboat no: 1, 2, 5, 6, 7, and

8). The captain's position does not surpass TLV because the captain's place is in front of the speedboat, and the distance between the captain and the engine is quite large. Especially in enclosed speedboats, the commotion generated by the machine will be automatically reduced by the walls surrounding the speedboat's body. However, the personnel will be directly exposed to the engine noise because they

are on a ship. Consequently, it is not remarkable that the noise level in this location exceeds the TLV and is higher.

The results of statistical analyses for each independent variable and dependent variable are displayed in Table 3. Consequently, almost all independent variables are associated with hearing impairments, with the exception of job position (p-value > 0.05).

Table 3. Correlation test results

Variable	Level	Hearing Status		Р-	OR	
varianie		Normal	Disturbances	Value	(95% CI)	
Noise	Average noise	19	27	*000	28.852	
	High noise	2	82	.000	(6.307 - 131.988)	
Age	≤ 37 years old	20	44	*000	29.545	
	> 37 years old	1	65	.000	(3.825 - 228.245)	
Job position	Skipper	1	19	.195		
	Speedboat crews	20	90	.193		
BMI	Normal	18	56	*800.	5.679	
	Abnormal	3	53	.000	(1.581 - 20.396)	
Smoking	No	17	2	*000	227.375	
	Yes	4	107	.000	(38.620 – <mark>1,338.683</mark>)	
Alcohol	No	19	24	*000	33.646 Mis	
	Yes	2	85	.000	(7.316 - 154.736)	
Working	≤ 7 years	20	47	*000	26.383	
experience	> 7 years	1	62	.000	(3.417 - 203.679)	
Work	1 - 5 hr	20	62	.002*	15.161	
duration	> 5 hr	1	47	.002	(1.964 - 117.047)	
Physical	Yes	14	7	*000	29.143	
Activities	No	7	102	.000	(8.890-95.534)	
Hobbies	Not related high sound	20	34	*000	44.118	
	Related high sound	1	75	.000	(5.686 - 342.303)	

^{*=} p < 0.05

Dis cussion

NIHL is defined as a prevalent occupational disease. It is an avoidable or preventable disease. The WHO (2020) identifies a number of additional noiserelated causes of hearing loss, including engine and detonation noise at work and excessive noise (18). According to the examination and analysis findings, there is a correlation between noise level and hearing loss. The noise level in the speedboat ranges from 81.3 to 92.1 dBA, with an average of 85.45 dBA (skipper's position) and 88.47 dBA at the crew's place during the trip. Based on Table 2, although the aggregate measurement results at the captain's position did not exceed the TLV, they were distinct from those at the crew's position. The crew from six speedboats, are subject to noise levels above the TLV. Another studies also found from the result of noise assessment among several kind of ships also greater than TLV (4,17,19-21) Article

The present study was found the sailors who exposed to noise levels above the average, have a 28 times greater likelihood of developing hearing loss. In the same similar results from prior research, Jumali et al. (2013) found that 34.85% of ferry boat's employees experienced hearing difficulties.

Even though hearing impairment is typically only experienced by employees with more than ten years working (22), in this study, 83.8% of

speedboat members were hearing-impaired despite an average length of service of only seven years. This is due to the fact that they are perpetually exposed to it every day, as well as several other factors. Zaw et al. (2020) mentioned that hearing loss can be caused by either a single exposure to exceedingly loud or prolonged exposure to loud noise (23). Noise-related pastimes also influence the incidence of hearing loss. Jannah et al. (2020) found that hobbies such as recreational exposure to loud noises from personal audio devices at high volume intensity and long periods also affect regular attendance at concerts, nightclubs, bars, and moderately noisy sporting events. The statistical findings of this study are consistent with Jannah et al. Aging is an additional cause, primarily due to the degeneration of sensory cells (24).

Age-related hearing loss was one of the most prevalent causes of high-frequency hearing loss, with its effects beginning around age forty (25,26). This may be due to the phenomenon of presbycusis, which is the progressive loss of hearing that occurs with advancing age (22,23). In this study, average crew's age is 37 years. The seafarers who have aged > 37 years and older were 29.545 times more likely to have hearing disturbances than those who were younger than 37. Similar results were also found in Zaw's study et al. (2020) workers aged > 35 were seven times more likely to suffer from hearing loss than those under 35 years (23).

Several modifiable lifestyle factors, including smoking, alcohol consumption (27), a high body mass index (BMI), and physical inactivity (28), have also been linked to hearing health. Statistical results also present the same as a lifestyle factors correlated with hearing impairment (p value < 0.05). Over fifty percent of speedboat members have a normal BMI, but the majority of have hearing loss (43%), as shown in Table 1. While other members with an abnormal BMI also experience hearing loss (40%). This study classifies abnormal BMI categories (≥ 25

kg/m²) as pre-obesity, class I obesity, and class II obesity. A recent study revealed that patients with a body mass index (BMI) greater than 27.5 kg/m² were 1.59 times more likely to experience sudden sensorineural hearing loss than those with a BMI less than 23.5 kg/m². A potential mechanism has been proposed to explain the association between adiposity and hearing loss. By stiffening or constricting the internal auditory artery, obesity-related atherosclerosis may reduce blood flow to the cochlea. Blood supply impairment to the cochlea may contribute to diminished auditory sensitivity (29).

Accelerometer-based comprehensive physical activity measurement measure weekly moderate-intensity physical activity (MPA) per participant by converting daily minutes to weekly minutes and each minute of vigorousintensity activity to two minutes of moderate-intensity exercise (30). However, in this study, physical activity was categorized based on whether or not respondents engaged in physical activities (such as walking, gym use, and other physical activities) in their free time. The study's results demonstrated physical activity correlates with hearing loss, as 102 respondents (79%) who did not engage in physical activity had hearing problems. Due to an increased likelihood of social isolation, individuals with moderate or severe hearing loss may engage in less physical activity than those with normal hearing. A person's inability to effectively monitor the acoustic environment could limit their capacity to engage in physical activities. Joint neural degeneration impacting the cochlear and vestibular sense organs (essential for balance) may also account for the association between hearing loss and physical activity (30).

Cigarette smoking has a substantial negative impact on the auditory system. Tobacco is known to impair hearing through direct and indirect mechanisms in the inner ear, resulting in hearing loss at various frequencies (31). The influence of

smoking on the blood circulation system in the cochlear organ is the cause of progressive high-frequency hearing loss, which occurs most frequently in old age (16). However, the exact mechanism smoking and NIHL remains linking unknown. Nicotine and other tobacco constituents may be ototoxic, causing harm to cochlear hair cells by increasing carbon monoxide hemoglobin or decreasing cochlear blood flow volume (32,33). Several studies discovered that the combined impact of smoking occupational noise was comparable to the sum of the effects of each factor individually (34).

this investigation, In alcohol divided consumption was categories: drinkers and non-drinkers. It was not specified how much alcohol was consumed daily by these alcoholics. The statistic results confirmed that alcohol consumption has a relationship with hearing loss. The speedboat crews have a 33.646 times the possibility experiencing hearing loss when drinking alcohol. Similar research conducted by Xu et al. (2021) showed that the prevalence of hearing impairment was significantly higher than those who never consumed alcohol (35). In contrast with Curhan et al. (2011) study, there was no association between alcohol consumed and the risk of hearing loss (36). Noise and alcohol consumption may have caused injury to the blood vessels, cochlear vestibular organs, and auditory nerves of the inner ear (37). The biological mechanisms underlying the synergistic effects of noise exposure and alcohol consumption on hearing loss are not fully understood, and further research is needed (35,37).

Conclusion

Overall, the results of noise measurement indicated that the captain position does not exceed the TLV. In contrast, six speedboats in the crew's place exceeded the threshold value based on the duration

of exposure. A total of 109 participants (83.8%) had hearing loss, with the majority having mild (40.8%), moderate (39.2%), and severe (3.8%) levels. Noise has an effect on hearing impairment, as shown by statistical results indicating that all variables in this study are related to hearing impairment, except for job position. Noise is not the only cause of because all variables are hearing loss interconnected. For instance, members of speedboats are susceptible to hearing loss due to prolonged exposure to noise, high decibel levels, and other factors. They will also experience hearing loss due to their age, BMI, and other lifestyle-related such as smoking, alcohol consumption, recreational activities, and daily exercise routines.

Limitation

Noise measurements were conducted only once with a specific route and speedboat model, despite speedboats operating daily. Therefore, the results of measurements cannot represent the entire sample. The purpose of the present study was to establish a correlation between the independent and dependent variables and not to identify the underlying cause. A Sound Level Meter was employed for measuring noise. Nonetheless, using a shoulder-mounted personal noise dosimeter that assesses an individual's exposure to noise over a single working will be more effective. Some questionnaires do not explicitly explain in detail the variables, only asking to consider general (consume or not consume; did or did not).

Acknowledgements

We thank all speedboat workers at the port of Tengkayu II. Tarakan City for their participated. This study was financially supported by Institute Community Service Act of Err Universitas Muhammadiyah le Err Kalimantan Timur

(Competitive research no: 670.9/LPPM/A.4/C/2021).

Conflict of interest No conflicts of interest have been declared by any of the authors.

References

- 1. Nelson DI, Nelson RY, Concha-Barrientos M, Fingerhut M The global burden of occupational noise-induced hearing loss. Am J Ind Med. 2005;48(6):446–58.
- Aliabadi M, Farhadian M, Darvishi E. Prediction of hearing loss among the noise-exposed workers in a steel factory using artificial intelligence approach. Int Arch Occup Environ Health [Internet]. 2015;88(6):779– 87. Available from: http://dx.doi.org/10.1007/s00420-014-1004-z
- 3. Febriyanto K, Suprayitno S. Implementation of Occupational Safety and Health for Using Sea Transportation (Case Study At Tengkayu Ii Port of Tarakan City). J Ilmu Kesehat. 2020;7(2):97–102.
- 4. Oldenburg M, Felten C, Hedtmann J, Jensen HJ. Physical influences on seafarers are different during their voyage episodes of port stay, river passage and sea passage: A maritime field study. PLoS One. 2020;15(4):1–12.
- Putland RL, de Jong CAF, Binnerts B, Farcas A, Merchant ND. Multisite validation of shipping noise maps using field measurements. Mar Pollut Bull [Internet]. 2022;179:113733. Available from: https://doi.org/10.1016/j.marpolbul. 2022.113733
- Esmaeili R, Zare S, Ghasemian F, Pourtaghi F, Saeidnia H, Pourtaghi G. Predicting and classifyng hearing loss in sailors working on speed vessels using neural networks: a field study. Med del Lav.

- 2022;113(3):1-13.
- WHO. 1 in 4 people projected to have hearing problems by 2050 [Internet]. Geneva; 2021. Available from: https://www.who.int/news/item/02-03-2021-who-1-in-4-peopleprojected-to-have-hearingproblems-by-2050
- Haile LM, Kamenov K, Briant PS, Orji AU, Steinmetz JD, Abdoli A, et al. Hearing loss prevalence and years lived with disability, 1990-2019: Findings from the Global Burden of Disease Study 2019. Lancet. 2021;397(10278):996–1009.
- 9. Brown CS, Emmett SD, Robler SK, Tucci DL. Global Hearing Loss Prevention. Otolaryngol Clin North Am. 2018;51(3):575–92.
- Albizu EJ, De Oliveira Gonçalves CG, De Lacerda ABM, Zeigelboim BS, Marques JM. Noise exposure and effects on hearing in Brazilian fishermen. Work. 2020;65(4):881– 9.
- Al-Dayyeni WS, Sun P, Qin J. Investigations of Auditory Filters Based Excitation Patterns for Assessment of Noise Induced Hearing Loss. Arch Acoust. 2018;43(3):477–86.
- 12. Muzaffar SJ, Orr L, Rickard RF, Coulson CJ, Irving RM. Mitigating Noise-Induced Hearing Loss After Blast Injury. Trauma (United Kingdom). 2019;21(2):121–7.
- 13. Moore BCJ. A Review of The Perceptual Effects of Hearing Loss for Frequencies Above 3 kHz. Int J Audiol. 2016;55(12):707–14.png Article
- Curovic L, Jeram S, Murovec J, Novakovic T, Rupnik K, Prezelj J. Impact of COVID-19 on environmental noise emitted from the port. Sci Total Environ. 2021;756(144147).
- 15. Notoatmodjo S. Metodologi Penelitian Kesehatan. Jakarta: Rineka Cipta; 2010.

- 16. Febriyanto K, Sunarti S, Ghozali, Rahman FF. Assessing Noise-Exposure and Daily Habits Can Cause Hearing Loss among Ladies Club at Nightclub. Assessing NoiseExposure and Daily Habits Can Cause Hearing Loss among Ladies Club at Nightclub. Indian J Public Heal Res Dev. 2019;3(10):853.
- 17. Somadayo S. Analisis Tingkat Kebisingan Mesin Speed Boat Ternate - Sofifi. J Sains, Sosisal dan Hum. 2022;2(1):74–82.
- 18. WHO. Deafness and hearing loss [Internet]. 2020 [cited 2023 Apr 20]. Available from: https://www.who.int/news-room/fact-sheets/detail/deafness-and-hearing-loss
- Levin JL, Curry WF, Shepherd S, Nalbone JT, Nonnenmann MW. Hearing loss and noise exposure among commercial fishermen in the gulf coast. J Occup Environ Med. 2016;58(3):306–13.
- Picu M, Picu L. Noise: a risk factor for sailors' health and navigation safety. Ann "Dunarea Jos" Univ Galati Fascicle II, Math Physics, Theor Mech. 2020;43(2):147–54.
- Schaal NC, Salaam R, Stevens M, Stubner A. Noise characterization of "effective quiet" areas on a U.S. Navy aircraft carrier. J Occup Environ Hyg [Internet].
 2019;16(5):329–35. Available from: https://doi.org/10.1080/15459624.20 19.1591628
- Jumali, Sumadi, Andriani S, Subhi M, Suprijanto D, Handayani WD, et al. Prevalence and Risk Factors Noise Induced Hearing Loss on the Ferry Machine Operator. Kesmas Natl Public Heal J. 2013;7(12):545.
- 23. Zaw AK, Myat AM, Thandar M,
 Htun YM, Aung TH, Tun KM, et al.
 Assessment of Noise Exposure and
 Hearing Loss Among Workers in
 Textile Mill (Thamine), Myanmar:

- A Cross-Sectional Study. Saf Health Work [Internet]. 2020;11(2):199–206. Available from: https://doi.org/10.1016/j.shaw.2020. 04.002
- 24. Jannah AR, Chairani A, Harjono Y. Risk Factors of Noised Induced Hearing Loss in Workers at Beveling Unit in Industry X, Bekasi, West Java. In: The 7th International Conference on Public Health. Solo; 2020. p. 188–96.
- Ashraf HD, Younus MA, Kumar P,
 Siddiqui MT, Ali SS, Siddiqui MI.
 Students 'Corner Frequency of hearing loss among textile industry workers of weaving unit in Karachi , Pakistan Abstract. J Pakistan Med Assoc. 2009;59(8):575–9.
- Ferrite S, Santana V. Joint effects of smoking, noise exposure and age on hearing loss. Occup Med (Chic III). 2005;55(1):48–53.
- Arumugam, Tukalan G, Anchery VA, Khosh A. Evaluation of Noise Induced Hearing Loss in Fishermen Who Work in Motor Boats in Karaikal. J Evol Med Dent Sci. 2015;4(73):12645–50.
- Curhan SG, Eavey R, Wang M, Stampfer M, Curhan GC. Body Mass Index, Waist Circumference, Physical Activity and Risk of Hearing Loss in Women. Am J Med. 2013;126(12):1–7.
- Hu H, Tomita K, Kuwahara K, Yamamoto M, Uehara A, Kochi T, et al. Obesity and risk of hearing loss: A prospective cohort study. Clin Nutr [Internet].
 2020;39(3):870-5. Available from: https://doi.org/10.1016/j.clnu.2019.0 3.020
- 30. Buchner DM. Promoting Physical Activity in Older Adults. Am Fam Physician. 2010;81(1):24–5.
- 31. Nandita G, Shivalingesh K,
 Anushtha K, Varnika S, Siddhant S,
 Shashank K. Effect of smoking on
 status of hearing: A cross-sectional

- study. Arch Otolaryngol Rhinol. 2022;8(3):009–13.
- 32. Fechter LD, Thorne PR, Nuttall AL.

 Effects of carbon monoxide on cochlear electrophysiology and blood flow. Hear Res.

 1987;27(1):37–45. Confused
- Browning GG, Gatehouse S, Lowe GDO. Blood Viscosity As a Factor in Sensorineural Hearing Impairment. Lancet. 1986;327(8473):121–3.
- 34. Pouryaghoub G, Mehrdad R,
 Mohammadi S. Interaction of
 smoking and occupational noise
 exposure on hearing loss: A crosssectional study. BMC Public Health.
 2007;7:3–7.
- 35. Xu Y, Liu J, Dong C, Guo S, Cao H,

- Zhang J, et al. Hearing status of low-income, middle-aged and elderly women in northern china: A population-based, cross-sectional roper Noun study. J Multidiscip Healthc. 2021;14(September):2617–24.
- Curhan SG, Eavey R, Shargorodsky J, Curhan GC. Prospective Study of Alcohol Use and Hearing Loss in Men. Ear Near, 2011;32(1):46–52.
- Men. Ear Near 2011;32(1):46–52.

 Qian P, Zhao Z, Liu S, Xin J, Liu Y, Hao Y, et al. Alcohol as a risk factor for hearing loss: A systematic review and meta-analysis. PLoS One [Internet]. 2023;18(1 January):1–13. Available from: http://dx.doi.org/10.1371/journal.po ne.0280641

Advanced Study_Occup Safety

ORIGINA	ALITY REPORT	
SIMILA	9% 16% 12% 5% ARITY INDEX INTERNET SOURCES PUBLICATIONS STUDENT	ΓPAPERS
PRIMAR	Y SOURCES	
1	www.ncbi.nlm.nih.gov Internet Source	2%
2	www.koreascience.or.kr Internet Source	2%
3	mattioli1885journals.com Internet Source	1 %
4	repository.unmul.ac.id Internet Source	1 %
5	www.researchgate.net Internet Source	1 %
6	media.neliti.com Internet Source	1 %
7	Gispen, Fiona E., David S. Chen, Dane J. Genther, and Frank R. Lin. "Association Between Hearing Impairment and Lower Levels of Physical Activity in Older Adults", Journal of the American Geriatrics Society, 2014. Publication	1 %

8	journals.umkt.ac.id Internet Source	1 %
9	journals.plos.org Internet Source	1 %
10	www.mpi.mb.ca Internet Source	1 %
11	link.springer.com Internet Source	1 %
12	Submitted to West Coast University Student Paper	1 %
13	academic.oup.com Internet Source	<1%
14	knepublishing.com Internet Source	<1%
15	research.library.mun.ca Internet Source	<1%
16	Xi Shen, Yingzhou Zhang, Tianqi Wang, Yuxin Sun. "Relational Database Watermarking for Data Tracing", 2020 International Conference on Cyber-Enabled Distributed Computing and Knowledge Discovery (CyberC), 2020 Publication	<1%
17	Submitted to Ngee Ann Polytechnic Student Paper	<1%

18	Aung K. Zaw, Aung M. Myat, Mya Thandar, Ye M. Htun, Than H. Aung, Kyaw M. Tun, Zaw M. Han. "Assessment of Noise Exposure and Hearing Loss Among Workers in a Textile Mill (Thamine), Myanmar: A Cross-Sectional Study", Safety and Health at Work, 2020 Publication	<1%
19	Submitted to Laredo Community College Student Paper	<1%
20	www.disability.illinois.edu Internet Source	<1%
21	Submitted to Southern New Hampshire University - Continuing Education Student Paper	<1%
22	Giorgio Burella, Lorenzo Moro. "A comparative study of the methods to assess occupational noise exposures of fish harvesters", Safety and Health at Work, 2020 Publication	<1%
23	documents.mx Internet Source	<1%
24	timesofindia.indiatimes.com Internet Source	<1%
25	www.dovepress.com Internet Source	<1%

Internet Source



Exclude quotes On Exclude bibliography On

Exclude matches

Off

Advanced Study_Occup Safety

PAGE 1

- ETS)
- **Sp.** This word is misspelled. Use a dictionary or spellchecker when you proofread your work.
- ETS)
- **Article Error** You may need to remove this article.
- ETS)
- **Verb** This verb may be incorrect. Proofread the sentence to make sure you have used the correct form of the verb.
- (ETS
- **Article Error** You may need to remove this article.
- (ETS)
- **Article Error** You may need to use an article before this word.
- (ETS
- **Article Error** You may need to use an article before this word.
- (ETS)
- **Article Error** You may need to use an article before this word. Consider using the article **the**.
- (ETS)
- **Article Error** You may need to use an article before this word. Consider using the article **the**.
- (ETS)
- Possessive This word may be a plural noun and may not need an apostrophe.
- (ETS)
- **Article Error** You may need to use an article before this word.

PAGE 2

- ETS)
- **Article Error** You may need to remove this article.
- (ETS)
- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.
- (ETS)
- **Wrong Article** You may have used the wrong article or pronoun. Proofread the sentence to make sure that the article or pronoun agrees with the word it describes.
- (ETS
- **Article Error** You may need to use an article before this word.
- **ETS**
- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.

- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice. P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice. **Article Error** You may need to use an article before this word. **Possessive** This word may be a plural noun and may not need an apostrophe. **Tone** This language may not be appropriate in an essay. **Article Error** You may need to remove this article. Missing "," You may need to place a comma after this word. PAGE 3 (ETS) Missing "," You may need to place a comma after this word. **Article Error** You may need to use an article before this word. to make sure that the article or pronoun agrees with the word it describes.
- Wrong Article You may have used the wrong article or pronoun. Proofread the sentence
- **Article Error** You may need to use an article before this word.
- **Possessive** This word may be a plural noun and may not need an apostrophe.
- **Article Error** You may need to use an article before this word.
- **Verb** This verb may be incorrect. Proofread the sentence to make sure you have used the correct form of the verb.
- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.
- **Article Error** You may need to use an article before this word.
- **Article Error** You may need to use an article before this word.



Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.



Article Error You may need to remove this article.



Confused You have used **personnel** in this sentence. You may need to use **personal** instead.



Frag. This sentence may be a fragment or may have incorrect punctuation. Proofread the sentence to be sure that it has correct punctuation and that it has an independent clause with a complete subject and predicate.



Missing "," You may need to place a comma after this word.

PAGE 5



Proofread This part of the sentence contains a grammatical error or misspelled word that makes your meaning unclear.



Prep. You may be using the wrong preposition.



Prep. You may be using the wrong preposition.



Article Error You may need to use an article before this word. Consider using the article **the**.



Missing "," You may need to place a comma after this word.



Tone This language may not be appropriate in an essay.



Tone This language may not be appropriate in an essay.



Article Error You may need to use an article before this word. Consider using the article **the**.



Missing "," You may need to place a comma after this word.



Wrong Article You may have used the wrong article or pronoun. Proofread the sentence to make sure that the article or pronoun agrees with the word it describes.



Confused You have used **A** in this sentence. You may need to use **an** instead.



Article Error You may need to use an article before this word.

Missing "," You may need to place a comma after this word.

Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your

(ETS

work.

- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.
- Article Error You may need to remove this article.
- Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.
- Article Error You may need to use an article before this word. Consider using the article the.
- Article Error You may need to use an article before this word. Consider using the article the.

PAGE 9

- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.
- P/V You have used the passive voice in this sentence. Depending upon what you wish to emphasize in the sentence, you may want to revise it using the active voice.
- Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.
- Missing "," You may need to place a comma after this word.
- Article Error You may need to use an article before this word.
- Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.
- Wrong Article You may have used the wrong article or pronoun. Proofread the sentence to make sure that the article or pronoun agrees with the word it describes.

PAGE 10

- **Prep.** You may be using the wrong preposition.
- Proper Noun If this word is a proper noun, you need to capitalize it.
- Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.





- Article Error You may need to use an article before this word.
- Article Error You may need to use an article before this word. Consider using the article the.
- Article Error You may need to use an article before this word. Consider using the article the.

PAGE 11

- **Proper Noun** If this word is a proper noun, you need to capitalize it.
- Article Error You may need to use an article before this word. Consider using the article the.
- **Confused** You have used **Hear** in this sentence. You may need to use **here** instead.
- S/V This subject and verb may not agree. Proofread the sentence to make sure the subject agrees with the verb.
- **Confused** You have used **Hear** in this sentence. You may need to use **here** instead.
- Sp. This word is misspelled. Use a dictionary or spellchecker when you proofread your work.