

NASKAH PUBLIKASI (MANUSCRIPT)

**POTENSI FRAKSI N-HEKSAN DAN N-BUTANOL DARI DAUN
KELUBUT (*Passiflora foetida* L.) TERHADAP PENGHAMBATAN
MONOMIKROBA BIOFILM *Pseudomonas aeruginosa* DAN
*Escherichia coli***

**POTENTIAL OF N-HEXANE AND N-BUTANOL FRACTIONS OF LEAF
KELUBUT (*Passiflora foetida* L.) ON BIOFILM INHIBITION OF
MONOMIRCOBIALS *Pseudomonas aeruginosa* AND *Escherichia coli***

Zufiha Citra Utami Masdar¹, Irmala Dewi², Chaerul Fadly Mochtar³



**DISUSUN OLEH :
ZUFIHA CITRA UTAMI MASDAR
1911102415129**

**PROGRAM STUDI S1 FARMASI
FAKULTAS FARMASI
UNIVERSITAS MUHAMMADIYAH KALIMANTAN TIMUR
2023**

Naskah Publikasi (Manuscript)

**Potensi Fraksi N-Heksan dari Daun Kelubut (*Passiflora Foetida L.*)
terhadap Penghambatan Monomikroba Biofilm *Pseudomonas
aeruginosa* dan *Escherichia coli***

**Potential of N-Hexane and N-Butanol Fractions of Leaf Kelubut
(*Passiflora foetida L.*) on Biofilm Inhibition of Monomircobials
Pseudomonas aeruginosa and *Escherichia coli***

Zufiha Citra Utami Masdar¹, Irmala Dewi², Chaerul Fadly Mochtar³



Disusun Oleh :

Zufiha Citra Utami Masdar

1911102415129

PROGRAM STUDI S1 FARMASI

FAKULTAS FARMASI

UNIVERSITAS MUHAMMADIYAH KALIMANTAN TIMUR

2023

LEMBAR PERSETUJUAN

**POTENSI FRAKSI N-HEKSAN DAN N-BUTANOL DARI DAUN
KELUBUT (*Passiflora foetida* L.) TERHADAP PENGHAMBATAN
MONOMIKROBA BIOFILM *Pseudomonas aeruginosa* dan *Escherichia
coli***

NASKAH PUBLIKASI

DISUSUN OLEH :

**Zufiha Citra Utami Masdar
1911102415129**

Pembimbing


Chaerul Fadly Mochtar Luthfi M. S.Farm., M.Biomed
NIDN. 11150992022

Mengetahui,

Koordinator Mata Ajar Skripsi


Apt. Rizki Nur Azmi, M.Farm
NIDN. 1102069201

LEMBAR PENGESAHAN

**POTENSI FRAKSI N-HEKSAN DAN N-BUTANOL DARI DAUN
KELUBUT (*Passiflora foetida* L.) TERHADAP PENGHAMBATAN
MONOMIKROBA BIOFILM *Pseudomonas aeruginosa* dan *Escherichia
coli***

NASKAH PUBLIKASI

DISUSUN OLEH :

**Zufiha Citra Utami Masdar
1911102415129**

Penguji 1



**Apt. Ika Ayu Mentari. M.Farm
NIDN. 1121019201**

Penguji 2




**Chaerul Fadly Mochtar Luthfi M. S.Farm., M.Biomed
NIDN. 11150992022**

Mengetahui,

Ketua Program Studi S1 Farmasi




**Apt. Ika Ayu Mentari. M.Farm
NIDN. 1121019201**

Potential of N-Hexane and N-Butanol Fractions of Leaf Kelubut (*Passiflora foetida L.*) on Biofilm Inhibition of Monomicrobials *Pseudomonas aeruginosa* and *Escherichia coli*

Zufiha Citra Utami Masdar.¹, Irmala Dewi.¹, Chaerul Fadly Mochtar^{1,2*}

¹Bachelor of Pharmacy, Faculty of Pharmacy, Universitas Muhammadiyah Kalimantan Timur, Indonesia

²Indonesian Biofilm Research Collaboration Center (IBRCC), Yogyakarta, Indonesia

Email : cfm782@umkt.ac.id

Abstract

Infectious diseases are disease conditions caused by pathogenic microorganisms. Infections caused by biofilms are a significant problem because the microbial community in the biofilm matrix is highly resistant to antimicrobial agents. Pseudomonas aeruginosa is one of the gram-negative opportunistic bacteria that is often involved in biofilm formation and Escherichia coli is also one of the bacteria that can cause infections in various human bodies such as in the digestive tract and urinary tract. Kelubut leaf is one of the herbs found in Indonesia and the Southeast Asian region. This plant has a number of bioactive components that have attracted attention and many have researched especially in the pharmaceutical field. Tests to determine biofilm inhibition were carried out by microdilution testing. The minimum biofilm inhibition concentration, which causes 50% inhibition (MBIC) was determined by crystal violet staining and read using a microplate reader at 620nm. Furthermore, it was analyzed using MBIC calculation and SPSS to evaluate the mechanism of effect. The results showed that the N-Hexane fraction and N-Butanol fraction showed biofilm inhibitory activity against P.Aeruginosa bacteria were 93.76%, 70.10% and on E.Coli were 85.36%, 72.01%. N-Hexane fraction can inhibit biofilm formation for 24 hours and has no difference with Chloramphenicol positive control, N-Butanol fraction can inhibit biofilm for 24 hours but there is a difference with Chloramphenicol control.

Keywords: Fraksi N-Heksan, Fraksi N-Butanol, Daun Kelubut, Biofilm, *Pseudomonas aeruginosa*, *Escherichia coli*

Introduction

According to a global study in 2019 there were 7.7 million deaths caused by infections. Infectious diseases are disease conditions caused by pathogenic microorganisms such as bacteria, viruses, fungi, or parasites. According to Kim et al., (2013), reported that about 70% of treatment failures are caused by fungal or bacterial polymicrobial infections that form biofilm. Biofilms are currently considered as mediators of infection with an estimated 80% of diseases associated with biofilm formation². Infections caused by biofilms are a significant problem because the microbial community within the biofilm matrix is highly resistant to antimicrobial agents. Microbes that form biofilms are usually resistant to common antimicrobial drugs and are able to evade host cells and the immune system, which can serve as a protective barrier. Biofilms are a factor of virulence and resistance as they proliferate as the clinical infection increases in the host cell. New antimicrobial agents are now increasingly in demand^{3,4}.

Pseudomonas aeruginosa is a type of opportunistic gram-negative bacteria that is often associated with biofilm development. Nosocomial infections (infections that occur in

healthcare facilities) are common, and it is believed that *Pseudomonas aeruginosa* is responsible for 10 to 20 percent of such infections. Biofilms on intravascular catheters or other implanted medical devices can increase the risk of infection and complicate therapy⁵. *Escherichia coli* is a type of bacteria that can cause infections in the urinary and gastrointestinal tracts, as well as other parts of the human body. A challenge in the treatment of *Escherichia coli* infections is antibiotic resistance, which can develop resistance to several types of antibiotics, making treatment difficult and requiring the use of stronger antibiotics or combinations of antibiotics⁶.

Bacterial infections caused by biofilms are difficult to treat. To kill bacteria in biofilm form requires 1000 times the dose of antimicrobials to achieve the same results as planktonic cells. Biofilms can be controlled by utilizing chemical compounds obtained from natural materials. Using chemical compounds as prevention and treatment of infection has become a common and effective approach in society and medical practice³.

Some herbs have been recognized to have antimicrobial properties that can be used as a substitute for formalin in some cases and are more easily available and more affordable. Kelubut plant with the Latin name *Passiflora foetida* L. is one of the herbs found in Indonesia and Southeast Asia. This plant has been used in traditional medicine by people in certain regions for a long time. Kelubut leaves have a number of bioactive components that have attracted attention and have been widely researched, especially in the pharmaceutical field.⁷

Material and Methods

Materials : Materials used were Kelubut Leaves from Samarinda, Kalimantan Timur. Other materials include the following: ethyl acetate, N-Hexane, N-Butanol, *Pseudomonas aeruginosa* and *Escherichia coli* bacteria, Blood Base Agar (BBA) media, Brain Heart Infusion (BHI) media, Aquadest, Crystal violet 1%, ethanol 96%, Nystatin, Chloramphenicol 1%, DMSO.

Apparatus: Analytical balance, blender, corong kaca, beaker glass, erlenmeyer, *rotary evaporator*, persolvent cup, *waterbath*, spatel, test tube, test tube rack, petri dish, ose needle, autoclave, bunsen, measuring cup, pipette, micropipette, blue tip, yellow tip, white tip, stirring rod, hotplate, incubator (If-2b), vortex, *microplate 96 wells* (Iwaki®), *microplate reader* (HiPo Biosan), *laminar air flow* (LAF), *microtube*.

Research Method Kelubut leaves were determined in Herbarium Mulawarman, Laboratory Ecology and Conservation Tropical Forest Biodiversity, Faculty of Forestry, Mulawarman University Samarinda with the document number 69/UN17.4.08/LL/2023.

Processing The plants were dry sorted and wet sorted and then dried by drying in the sun. After the kelubut leaves are dry, they are pulverized into powder (*simplisia*) with a blender and stored in a tightly closed jar⁸.

Extraction Kelubut leaves were extracted with ethyl acetate by maceration method in a ratio 1:3. Maceration was carried out for 15 days at a constant temperature and periodic stirring was done. Every five days the macerate was filtered with a cloth to separate it from the pulp. After that, the extract will be concentrated using a *vacuum rotary evaporator* at 55°C to get a thick extract, and then the yield is calculated.⁹

Fractionation The result of thick ethyl acetate extract, measure as much as 5 grams. Then mix it with 100 mL of boiling water in an erlenmeyer flask. After the contents of the mixing container are put into a separatory funnel, 100 mL of N-Hexane is added and the mixture is stirred for 1 minute. Let stand for a while after shaking to produce two different stages. Fractionation was carried out repeatedly with the addition of N-Hexane until the fraction obtained changed color to clear. The resulting fraction is concentrated until it becomes a thick extract. Then the same was done for fractionation

using N-Butanol solvent.¹⁰

Microbial Preparation *Pseudomonas aeruginosa* and *Escherichia coli* bacteria were cultured in BBA media and incubated at 37°C for 24 hours. Then, *Pseudomonas aeruginosa* and *Escherichia coli* bacteria were re-cultured into liquid media, namely BHI and incubated at 37°C for 24 hours. The optical density (OD₆₀₀) of the microbial cultures was adjusted to the standard of (0,5 Mc Farland ~1,5 x 10⁸ CFU/ml)³.

Biofilm Inhibition The microdilution technique was used to test biofilm inhibition. A 96-well polystyrene microtiter plate with a flat bottom was used for the test, and concentration levels of 25%, 50%, and 100% were used. 100µL of *Pseudomonas aeruginosa* and *Escherichia coli* bacterial suspensions were put into the microplate wells and 100µL of N-Hexan & N-Butanol fraction solutions of kelubut leaves with concentrations (25%, 50%, 100%) were added. Furthermore, as a control medium, a bacterial suspension was given without any microbial growth, and as a growth control, a bacterial suspension was used. To compare the test findings, a microbiological culture containing 100 µL of chloramphenicol was used as a positive control in the wells. After that, it was cultured for 24 hours at 37°C in an incubator. To remove residual water, the microplate was then dried at room temperature and cleaned three times using distilled water. To color the growing biofilm, 125 µL of 1% crystal violet solution was applied to each well. For fifteen minutes, the microplate was incubated at room temperature. After the incubation period, each well received 200 µL of 96% ethanol, and the microplate was thoroughly cleaned three times using distilled water to remove residual crystal violet¹¹. The use of crystal violet as a dye increases precision and enables quantitative detection of biofilms. The basic dye called crystal violet binds to polysaccharides and negatively charged compounds in the extracellular matrix. As a result, the biofilm matrix and living and dead cells can be stained with crystal violet¹¹. The reading of biofilm degradation results was carried out using a 620nm Optical Density (OD) microplate reader. The OD value is then used to calculate the biofilm inhibition^{3,12}.

Data Analysis

The percentage of inhibition and degeneration concentration was counted using the formula below:

$$\% \text{ Inhibition} = \frac{(OD \text{ growth control} - OD \text{ sample})}{OD \text{ growth control}} \times 100\%$$

Statistical Analysis Analysis using the SPSS for Windows program, *One way Analysis of Variance* (ANOVA) test was used to test the research data. With a data significance level of 0.05, this ANOVA is a parametric test that compares the average differences of two or more treatment groups using a numerical data scale. The *Shapiro Wilk* test is used to determine whether the data distribution is normal. Data is said to be normal if a significance value greater than 0.05 is found. While the data is not normally distributed if <0.05. The Lavene test was used in the sample homogeneity test, and the findings that showed homogeneous data had a significance value >0.05¹³. The *Kruskal Wallis* test will be used to assess the data of this study if it does not meet the requirements of the one-way ANOVA test. After using one-way ANOVA, *Tukey's post hoc* test was conducted. However, if *Kruskal Wallis* is used in data analysis, then proceed to the Man Withney test¹⁴.

Results and Discussion

Kelubut leaves (*Passiflora foetida* L.) that have been determined are dried then mashed and macerated extraction is carried out as much as 1000 grams for 5 days

using 96% ethyl acetate solvent in a ratio of 1: 3 and repeated 3 times. Extraction of dried leaves (1000gram) produced 25.58 grams of ethyl acetate extract (13.04%). Then the liquid-liquid fractionation process was carried out with N-Hexan solvent which obtained a thick extract of 13.04 grams and N-Butanol as much as 5.17 gram. Biofilms caused by bacterial and fungal infections are one of the health problems that about 70% result in treatment failure. Infections caused by biofilms are a significant problem because the microbial community within the biofilm matrix is highly resistant to antimicrobial agents. Microbes that form biofilms are usually resistant to common antimicrobial drugs and are able to evade host cells and the immune system, which can serve as a protective barrier. In this study, we evaluated the potential of N-Hexane and N-Butanol fractions of kelubut leaves in the inhibition of monomicrobial biofilm formation of *Pseudomonas aeruginosa* and *Escherichia coli*. The results showed that the N-Hexane and N-Butanol fractions of kelubut leaves could inhibit 50% of biofilm formation (Figure 1 and Figure 2).

The concentration of chloramphenicol used in this study was 1%. Chloramphenicol is an antibiotic that is useful against a wide variety of bacteria. Chloramphenicol is a bacteriostatic antibiotic that at high enough concentrations can also kill bacteria. Its function is to block the formation of peptide bonds by attaching to ribosomes, preventing protein synthesis¹⁵. The use of positive controls has the aim of comparing the results of inhibition of biofilm formation^{15,16}.




Fig. 1: Biofilm inhibition activity monomicrobial *Pseudomonas aeruginosa*

In this result, it was reported that the N-Hexane fraction of kelubut leaves provided greater biofilm formation inhibitory activity compared to the N-Butanol fraction. The N-Hexane fraction gave an activity of 93.76% while the N-Butanol fraction gave an activity of 70.10% at 100% concentration. The activity of the N-Hexane fraction was almost the same as the positive control, namely chloramphenicol 1% at 95.27% (Figure 1). These results indicate that concentrations of 100% and 50% of the N-Hexane fraction and N-Butanol fraction of kelubut leaves are able to inhibit the formation of *Pseudomonas aeruginosa* monomicrobial biofilms above 50%.




Fig. 2: Biofilm Inhibition activity monomicrobial *Escherichia coli*

The results of the study in Figure 2 show that the 100% N-Hexan fraction provides biofilm formation inhibitory activity against *Escherichia coli* monomicrobials of 85.36% and at 100% N-Butanol fraction concentration provides activity of 72.01%. At a concentration of 25% N-hexan fraction has a biofilm formation inhibitory activity of 52.48% and at 25% N-Butanol fraction of 55.95%.

Biofilm inhibition mechanism and planktonic inhibition mechanism are different. In biofilms, bacteria together form a group or community so as to produce a more complex and stronger defense, while in planktonic bacteria are only a single cell and live freely so that antimicrobial agents can cause more damage naturally to defense cells and reach target cells¹⁷.

The results of the biofilm inhibition test presentation above show an increase in the percentage of biofilm inhibition against *Pseudomonas aeruginosa* and *Escherichia coli* bacteria along with an increase in the concentration used. This means that the higher the concentration used, the higher the biofilm inhibition produced. An increase in the potential inhibition of biofilm formation was found at a concentration of 100% N-Hexan fraction against *Pseudomonas aeruginosa*, there was no significant difference ($p > 0.05$) when compared to the positive control, which means there is no significant difference in the inhibition of biofilm formation. At a concentration of 100%, the N-Butanol fraction found a significant difference ($p < 0.05$) when compared to the positive control, which means it has a difference in inhibiting the formation of *Pseudomonas aeruginosa* biofilm. At a concentration of 100% N-Hexan fraction against *Escherichia coli* when compared to the positive control did not have a significant difference ($P > 0.05$) which means it does not have a meaningful difference. At a concentration of 100%, the N-Butanol fraction has a significant difference ($p < 0.05$) when compared to the positive control, which means it has a significant difference.

According to Miquel et al., (2016) Plants have bioactive chemicals that can inhibit the process of bacterial attachment to solid surfaces and the formation of extracellular polymer (EPS) scaffolds. This may result in anti-biofilm action. Kelubut leaves contain alkaloids, steroids, tannins, saponins, coumarins, tyrosine, glycine, and flavonoids. However, in this study, N-Hexan and N-Butanol fractionation was carried out so it is possible that the compounds of interest in this N-Hexan fractionation are steroids/triterpenoids and saponins. Steroids/triterpenoids prevent the production of new proteins that accumulate and modify the constituent elements of bacterial cells, thus inhibiting bacterial development¹⁹. By attaching to the bacterial biofilm layer, saponins work by reducing the number of bacterial cells thereby disrupting their permeability and making their cell walls brittle and eventually resulting in death²⁰. N-

Butanol fractionation allows the attraction of polar compounds, namely alkaloids, flavonoids, and tannins. Alkaloids have the ability to inhibit the synthesis of peptidoglycan in bacterial cells thus inhibiting the formation of the cell wall layer properly, while flavonoids can interfere with energy transduction in the bacterial cytoplasmic membrane, inhibiting the production of ATP which is essential for bacterial life. Flavonoids can also inhibit bacterial motility by inhibiting flagellum synthesis and inhibiting enzymes involved in bacterial movement, and tannin compounds can cause cell damage by denaturing proteins²¹.

Conclusion

In conclusion, the available data show that the N-Hexan fraction and N-Butanol fraction of kelubut leaves (*Passiflora foetida* L.) have inhibitory activity onomicrobial biofilm formation *Pseudomonas aeruginosa* and *Escherichia coli*. N-Hexan fraction is the most active fraction as inhibition of biofilm formation with the highest percentage of inhibition against *Pseudomonas aeruginosa* 93.76% and against *Escherichia coli* 85.36%. The N-Hexan fraction when compared with the positive control chloramphenicol 1% did not have a significant difference ($p>0.05$) in inhibiting biofilm formation of *Pseudomonas aeruginosa* and *Escherichia coli* monomicrobials. The N-Butanol fraction also has inhibitory activity of biofilm formation with a percent inhibition of 70.10% on *Pseudomonas aeruginosa* bacteria and a percent inhibition of 72.01% on *Escherichia coli* bacteria. However, the N-Butanol fraction has a significant difference when compared to the positive control of chloramphenicol 1%.

Acknowledgement

We thank the Indonesian Biofilm Research Collaboration Center BRIN-UGM-Unsoed-UMKT for providing bacterial isolates and some consumables.

References

1. Kim, S. H., Yoon, Y. K., Kim, M. J. & Sohn, J. W. Risk factors for and clinical implications of mixed *Candida*/bacterial bloodstream infections. *Clin. Microbiol. Infect.* **19**, 62–68 (2013).
2. Anggraini, W., Purwanto, D. A., Isnaeni, Kusumawati, I. & Suryanto. a Systematic Review of Potential Phytochemical Compound Bark of *Parameria Laevigata* on Biofilm Formation. *Int. J. Appl. Pharm.* **14**, 21–26 (2022).
3. Hamzah, H., Hertiani, T., Pratiwi, S. U. T., Nuryastuti, T. & Gani, A. P. Antibiofilm studies of zerumbone against polymicrobial biofilms of *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Candida albicans*. *Int. J. Pharm. Res.* **12**, 1307–1314 (2020).
4. Khan, F., Tabassum, N., Jeong, G. J., Jung, W. K. & Kim, Y. M. Inhibition of Mixed Biofilms of *Candida albicans* and *Staphylococcus aureus* by β -Caryophyllene-Gold Nanoparticles. *Antibiotics* **12**, 1–17 (2023).
5. Gunardi, W. D. Mekanisme Biomolekuler *Pseudomonas aeruginosa* dalam Pembentukan Biofilm dan Sifat Resistensi terhadap Antibiotika. *J. Kedokt. Meditek* **22**, 1–7 (2017).
6. Wahyudi, D. & Wael, S. Pengaruh Ekstrak *Alpinia galanga* L Terhadap Produksi Biofilm pada *Escherichia coli*. *J. Farm. (Journal Pharmacy)* **10**, 24–30 (2021).
7. Jufri, N., Laga, A. & Faidah, A. N. EFEKTIVITAS EKSTRAK RAMBUSA (*Passiflora foetida* L.) DALAM MENGHAMBAT BAKTERI, KHAMIR DAN PENGARUHNYA PADA TOTAL MIKROBA TAHU *Makal. yang disajikan pada ...* (2022).
8. Kining, E., Falah, S. & Nurhidayat, N. The *In Vitro* Antibiofilm Activity of Water Leaf Extract of Papaya (*Carica papaya* L.) against *Pseudomonas aeruginosa*.

- Curr. Biochem.* **2**, 150–163 (2017).
9. Mukhriani, Tahar, N. & Astha, A. S. W. Uji Aktivitas Antibakteri Hasil Fraksinasi Dari Ekstrak Metanol Daun Katuk (*Sauropus androgynus*) Terhadap Beberapa Bakteri Patogen. *Jf Fik Unam* **12**, 12 (2014).
 10. Irvan herdiana, N. A. Jurnal Ilmiah Kesehatan 2022 Jurnal Ilmiah Kesehatan 2022. *J. Ilm. Kesehat.* **21**, 14–19 (2020).
 11. Dewi, Z. Y., Nur, A. & Hertriani, T. Efek antibakteri dan penghambatan biofilm ekstrak sereh (*Cymbopogon nardus* L.) terhadap bakteri *Streptococcus mutans*. *Maj. Kedokt. Gigi Indones.* **20**, 136 (2015).
 12. Hamzah, H., Hertiani, T., Utami Tunjung Pratiwi, S. & Nuryastuti, T. The Inhibition Activity of Tannin on the Formation of Mono-Species and Polymicrobial Biofilm *Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Candida albicans*. *Maj. Obat Tradis.* **24**, 110 (2019).
 13. Abidah, H. Y. Uji aktivitas antibiofilm ekstrak daun murbei hitam (*Morus nigra* L.) terhadap biofilm *Escherichia coli*. *Skripsi. Fak. Kedokt. dan Ilmu Kesehat. Univ. Islam Negeri Maulana Malik Ibrahim Malang* 1–160 (2020).
 14. Aviantina, M. E. Uji Aktivitas Penghambatan Biofilm Ekstrak Etanol Daun Kirinyu (*Chromolaena odorata* (L.) R. M. King & H. Rob.) terhadap Bakteri *Staphylococcus aureus*. 11–12 (2019).
 15. Dian, R., . F. & Budiarso, F. Uji Resistensi Bakteri *Escherichia Coli* Yang Diisolasi Dari Plak Gigi Terhadap Merkuri Dan Antibiotik Kloramfenikol. *J. e-Biomedik* **3**, (2015).
 16. Anggraini, W., Nisa, S. C., Da, R. R. & Ma, B. Aktivitas antibakteri ekstrak etanol 96 % buah blewah (*cucumis melo* l . Var . *Antibacterial activity of 96 % ethanol extract cantaloupe fruit (cucumis melo* l . Var . *Cantalupensis*) against *escherichia coli* bacteria. *Pharm. J. Indones.* **5**, 61–66 (2019).
 17. Galié, S., García-Gutiérrez, C., Miguélez, E. M., Villar, C. J. & Lombó, F. Biofilms in the food industry: Health aspects and control methods. *Front. Microbiol.* **9**, 1–18 (2018).
 18. Miquel, S., Lagrèfeuille, R., Souweine, B. & Forestier, C. Anti-biofilm activity as a health issue. *Front. Microbiol.* **7**, 1–14 (2016).
 19. Noviyanti, P. S. & Tarigan, D. Uji FITOKIMIA, TOKSISITAS DAN AKTIVITAS ANTIBAKTERI TERHADAP EKSTRAK ETANOL DAUN RAMBUSA (*Passiflora foetida* L.) TERHADAP BAKTERI *Staphylococcus aureus* DAN *Escherichia coli*. *J. Kim. Mulawarman* **12**, 31–36 (2014).
 20. Martin Widayat, M., Purwanto & Dewi, A. S. P. Daya Antibakteri Infusa Kulit Manggis (*Garcinia mangostana* L) terhadap *Streptococcus mutans* (Antibacterial of Mangosteen peel infuse (*Garcinia mangostana* L) Against *Streptococcus mutans*). *e-Jurnal Pustaka Kesehat.* **4**, 514–518 (2016).
 21. Lestari, D. R. S., Soegianto, L. & Hermanu, L. S. Potensi Antibakteri dan Antibiofilm Ekstrak Etanol Bunga Bintaro (*Cerbera odollam*) terhadap *Staphylococcus aureus* ATCC 6538. *J Pharm. Sci.* **4**, 30–35 (2017).



SURAT KETERANGAN ARTIKEL PUBLIKASI

Assalamu'alaikum Warahmatullahi wbarakatuh

Saya yang bertanda tangan dibawah ini:

| | | |
|---------------|---|-----------------------------------|
| Nama | : | Chaerul Fadly Mochtar L, M.Biomed |
| NIDN | : | 115099202 |
| Nama | : | Zufiha Citra Utami Masdar |
| NIM | : | 1911102415129 |
| Fakultas | : | Farmasi |
| Program Studi | : | S1 Farmasi |

Menyatakan bahwa "Aktivitas Fraksi N-Heksan dari Daun Kelubut (*Passiflora foetida* L) Terhadap Penghambatan Polimikroba Biofilm *Pseudomonas aeruginosa* dan *Escherichia coli*"

Demikian surat keterangan ini dibuat untuk dapat dipergunakan sebagaimana mestinya.

Wassalamu'alaikum Warahmatullahi wbarakatuh

Samarinda, Selasa 9 Januari 2024

Mahasiswa/i

Zufiha Citra Utami Masdar
NIM. 1911102415129

Dosen Pembimbing Skripsi

Chaerul Fadly Mochtar L, M. Biomed
NIDN. 115099202

| | | | |
|--|--|-----------------------|---------------------|
| International Journal of Agricultural Sciences and Veterinary Medicine | ACTIONS OF TROPICAL FRUIT ANNONA MURICATA LINN | | |
| International Research Journal for Quality in Education | 4182 Principle Component Analysis of Ascorbic acid ameliorating effects in salt-stressed tomato germplasm at the seedling stage | 12/1/2023 2:28:33 AM | Pinjari Osman Basha |
| NanoMatChemBioDev | 4183 Green Synthesis of Silver Nanoparticles from Brown Seaweed <i>Padina gymnospora</i> . | 12/1/2023 2:42:54 AM | L Archana |
| Yoga and Spiritualism Journal | 4184 Detection and Recurrence of Breast Cancer through Image Processing and Attention Awareness : A Comparative Analysis of Algorithms | 12/2/2023 1:47:58 AM | Alan George |
| Global Research in Medical Sciences | 4185 Current Trends in Nutraceuticals: Health benefits and Marketed formulations | 12/2/2023 3:12:45 AM | Ittishree |
| Research Journal of Bioinformatics | 4186 Current Trends in Nutraceuticals: Health benefits and Marketed formulations | 12/2/2023 3:12:50 AM | Ittishree |
| | 4187 Current Trends in Nutraceuticals: Health benefits and Marketed formulations | 12/2/2023 3:12:54 AM | Ittishree |
| | 4188 A study exploring the optimization of nutritional parameters of bioprocesses to maximize the production of keratinase by <i>Bacillus subtilis</i> RAS04. | 12/2/2023 10:44:11 PM | Rishi A Soni |
| | 4189 RIP, State, Production, Pharmacology, and Perspectives in medicinal plants (<i>Bryonia</i>) – A review | 12/3/2023 10:33:33 PM | Sahu Hemshankar |
| | 4190 Potential Probiotic Bacteria <i>Lactobacillus kefir</i> YK4 from Indonesian Kefir Grain as a Producer of Health Promoting Bioactive Compounds Identified by Nano-LCMS and In Silico | 12/3/2023 11:09:02 PM | Dandy Yusuf |
| | 4191 Comparison of cultivated <i>Cordyceps militaris</i> and wild <i>Ophiocordyceps sinensis</i> using high-performance thin-layer chromatography | 12/4/2023 10:41:41 PM | Alok Malaviya |
| | 4192 Potential of N-Hexane and N-Butanol Fractions of Leaf <i>Kelubut</i> (<i>Passiflora foetida</i> L.) on Biofilm Inhibition of Monomicrobial <i>Pseudomonas aeruginosa</i> and <i>Escherichia coli</i> | 12/7/2023 9:13:10 AM | Mochtar |

UGC Approved Journal
(Important for Indian Authors):
**Research Journal of
Biotechnology**



Publisher:
Dr. Jyoti Garg
Address:
Sector AG /So, Vihav Nagar.

<http://worldresearchersassociations.com/biotech.aspx>